

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963 7 A

Kacasasan aastassa saasassa saasaasan baasaasan kacaasaaan eessississa l

Department of Mathematics and Statistics The University of South Carolina Columbia, South Carolina 29208

AFOSR TR- 85-0865



AD-A160 302


ON THE ASYMPTOTIC PROPERTIES OF A KERNEL-TYPE QUANTILE ESTIMATOR FROM CENSORED SAMPLES

by

Y. L. Lio, W. J. Padgett, and K. F. Yu

University of South Carolina Statistics Technical Report No. 104 62G05-13





OTIC FILE COPY

85 10 11 119

ON THE ASYMPTOTIC PROPERTIES OF A KERNEL-TYPE QUANTILE ESTIMATOR FROM CENSORED SAMPLES *

Ъу

Y. L. Lio, W. J. Padgett, and K. F. Yu

University of South Carolina Statistics Technical Report No. 104 62G05-13

May, 1985

Department of Mathematics and Statistics University of South Carolina Columbia, South Carolina 29208 OCT 1 5 1985

Research supported by the U.S. Air Force Office of Scientific Research and Army Research Office under grant AFOSR-84-0156.

** Partially supported by a University of South Carolina Research and Productive Scholarship Grant.

Abstract

Some asymptotic results for a kernel type estimator of the quantile function from right-censored data are obtained. The estimator is defined by $Q_n(p) = h_n^{-1} \int_0^1 \hat{Q}_n(t) K((t-p)/h_n) dt$, which is smoother than the usual product-limit quantile function $\hat{Q}_n(p) = \inf\{t: \hat{F}_n(t) \ge p\}$, where \hat{F}_n denotes the product-limit estimator of the lifetime distribution F_0 . Under the random censorship model and general conditions on h_n , K, and F_0 , the asymptotic normality of $Q_n(p)$ is proven. In addition, an approximation to Q_n is shown to be asymptotically uniformly equivalent to Q_n in mean square.

Acces	sion For	
DTIC Unann	GRA&I TAB Ounced fication	
By	ibution/	
Avai	lability Codes	DTIG
Dist	Avail and/or Special	CCMS IN SPECIAL 23
A-1		

<u>Key Words</u>: Random right-censorship; Kernel estimation; Product-limit quantile function; Asymptotic normality; Mean-square convergence.

1. Introduction

In reliability and medical studies, it is often of interest to estimate various quantiles of the unknown lifetime distribution. In particular, the median lifetime and extreme quantiles are of interest to the experimenter in such studies. In many life testing and medical follow-up experiments, however, arbitrarily right-censored data arise, and it is important to be able to estimate the quantiles of interest based on the censored data. For such data, some kernel-type quantile estimators are considered in this paper which give smoother estimates than the usual product-limit quantile function.

For any probability distribution function G, denote the quantile function by $Q(p) \equiv G^{-1}(p) = \inf\{x \colon G(x) \geq p\}$, $0 \leq p \leq 1$. For a random (uncensored) sample Y_1, \ldots, Y_n from G, the sample quantile function $G_n^{-1}(p) = \inf\{x \colon G_n(x) \geq p\}$, $0 \leq p \leq 1$, has been used to estimate Q(p), where G_n denotes the sample distribution function. Cs0rg& (1983) gave many of the known results concerning $G_n^{-1}(p)$. Also, Falk (1984) studied the relative deficiency of the sample quantile with respect to kernel-type estimators, and Falk (1985) obtained asymptotic normality for kernel estimators. Yang (1985) has obtained some convergence properties of kernel estimators of Q(p) and gave some simulation results comparing kernel-type estimators with other estimators. For arbitrarily right-censored data, Sander (1975) proposed estimation of Q(p) by the quantile function of the product-limit estimator, and she and Cheng (1981) derived some asymptotic properties of that estimator. Also, Cs0rg& (1983) presented strong approximation results for that estimator.

Recently, Padgett (1985) studied a smoothed nonparametric estimator of

Q(p) from arbitrarily right-censored data based on the kernel method. It was shown that his estimator, mentioned briefly by Parzen (1979, p. 119), was strongly consistent, and a small Monte Carlo study was performed to compare the estimator with the product-limit estimator. In addition, a simple approximation to this kernel estimator was shown to be almost surely asymptotically equivalent to it.

The purpose of this paper is to further study the asymptotic properties of the estimators proposed by Padgett (1985). In particular, the asymptotic normality will be proven, and the asymptotic equivalence in mean square of the estimator and its approximation will be shown under general conditions on the kernel function, bandwidth sequence, lifetime distribution, and censoring mechanism.

2. Notation and Preliminaries

Let $X_1^0, X_2^0, \ldots, X_n^0$ denote the true survival times of n items or individuals which are censored on the right by a sequence U_1, U_2, \ldots, U_n , which in general may be either constants or random variables. It is assumed that the X_1^0 's are nonnegative independent identically distributed random variables with common unknown distribution function F_0 and unknown quantile function

$$Q^{0}(p) \equiv \xi_{p}^{0} = \inf\{t: F_{0}(t) \ge p\}, 0 \le p \le 1.$$

The observed right-censored data are denoted by the pairs (X_i, Δ_i) , i=1,...,n, where

$$X_{i} = \min\{X_{i}^{o}, U_{i}^{o}\}, \quad \Delta_{i} = \begin{cases} 1 & \text{if } X_{i}^{o} \leq U_{i} \\ 0 & \text{if } X_{i}^{o} > U_{i} \end{cases}.$$

For the asymptotic results of this paper, the random right-censorship model will be assumed, that is, V_1, \ldots, V_n constitute a random sample from a distribution H (usually unknown) and are independent of X_1^0, \ldots, X_n^0 . The distribution function of each X_1 , i=1,...,n, is then $F = 1 - (1-F_0)(1-H)$.

A popular estimator of the survival function $S_O(t) = 1 - F_O(t)$ based on (X_i, Δ_i) , i=1,...,n, is the product-limit estimator, proposed by Kaplan and Meier (1958) as the "nonparametric maximum likelihood estimator." Let (Z_i, Δ_i') , i=1,...,n, denote the ordered X_i 's along with their corresponding Δ_i 's. The product-limit estimator of $S_O(t)$, shown to be "self-consistent" by Efron (1967), is defined by

$$\hat{P}_{n}(t) = \begin{cases} 1, & 0 \le t \le Z_{1}, \\ \frac{k-1}{n} & (\frac{n-i}{n-i+1})^{\Delta_{i}^{t}}, & Z_{k-1} < t \le Z_{k}, & k=2,...,n \\ 0, & t > Z_{n}. \end{cases}$$

Denote the product-limit estimator of $F_0(t)$ by $\hat{F}_n(t) = 1 - \hat{P}_n(t)$, and let s_j denote the jump of \hat{P}_n at Z_j , that is

$$s_{j} = \begin{cases} 1 - \hat{P}_{n}(z_{2}), & j = 1 \\ \hat{P}_{n}(z_{j}) - \hat{P}_{n}(z_{j+1}), & j = 2, ..., n-1 \\ \hat{P}_{n}(z_{n}), & j = n. \end{cases}$$

Note that $s_j=0$ if and only if $\Delta_j^i=0$, j< n, i.e. whenever Z_j is a censored observation. Also, denote $S_i = \hat{F}_n(Z_{i+1}) = \sum_{j=1}^{r} s_j$, $i=1,\ldots,n$, with $S_0 = 0$, $Z_0 = 0$, and $Z_{n+1} = Z_n + \varepsilon$, for some positive constant ε . It is natural to estimate ξ_p^0 by the product-limit quantile function $\hat{Q}_n(p) = \hat{\xi}_p^0 = \inf\{t: \hat{F}_n(t) \ge p\}$. Sander (1975) and Cheng (1981) obtained

asymptotic normality results for $\hat{\xi}_p^o$. Padgett (1985) smoothed \hat{Q}_n^o by the kernel method to obtain the estimator

$$Q_{n}(p) = h_{n}^{-1} \int_{0}^{1} \hat{Q}_{n}(t) K((t-p)/h_{n}) dt$$

$$= h_{n}^{-1} \sum_{i=1}^{n} Z_{i} \int_{S_{i-1}}^{S_{i}} K((t-p)/h_{n}) dt, \qquad (2.1)$$

where K is an appropriate kernel function and $\{h_n\}$ is a bandwidth sequence. Also, a simpler kernel-type estimator which is an approximation to (2.1) was defined by

$$Q_n^*(p) = h_n^{-1} \sum_{i=1}^n Z_i^s K((S_i - p)/h_n).$$
 (2.2)

Note that only the uncensored observations actually appear in the sums of (2.1) and (2.2).

In the next section, the asymptotic normality of $Q_n(p)$ and $Q_n^*(p)$ will be obtained. The following general conditions on the kernel function, the bandwidth sequence, and the lifetime and censoring distributions will be assumed:

- (h.1) $h_n \to 0$ as $n \to \infty$;
- (K.1) K(x) is a bounded probability density function which has finite support, i.e. K(x) = 0 for |x| > c for some c > 0;
- (K.2) K is symmetric about zero;

$$|K(x) - K(y)| \le \Gamma |x - y|$$
;

- (F.1) $\mathbf{f}_{\mathbf{o}}$ is continuous with density function $\mathbf{f}_{\mathbf{o}}$; and
- (F.2) $H(T_{\overline{F}_0}) < 1$, where $T_{\overline{F}_0} = \sup\{t: F_0(t) < 1\}$.

It should be noted that these conditions are not prohibitive and (F.1) and (F.2) are similar to conditions required by Cheng (1981). Also, (F.2) insures that observations will be available from the entire support of $\mathbf{F_0}$, a common condition in random right-censorship models.

3. The Main Results

In this section, the main results are summarized in Theorems 3.1 and 3.2. The proofs will be presented in the next section. Theorem 3.1 gives conditions for the asymptotic normality of $Q_n(p)$. The asymptotic uniform mean-squared equivalence of Q_n and Q_n^* will be shown in Theorem 3.2.

Theorem 3.1. Assume, in addition to conditions (h.1), (K.1) - (K.3), (F.1), and (F.2), that the derivative f'_0 is continuous at ξ^0_p and $f_0(\xi^0_p) > 0$. Suppose $\{h_n\}$ is such that $n^{1/4}h_n + 0$ as $n \to \infty$. Then for 0 , where <math>T < 1, as $n \to \infty$, $\sqrt{n}[Q_n(p) - Q^0(p)] \to Z$ in distribution, where Z is a normally distributed random variable with mean zero and variance

$$\sigma_{\rm p}^2 = (1 - {\rm p})^2 \int_0^{\xi_{\rm p}^0} [1 - {\rm F}({\rm u})]^{-2} \frac{{\rm d} F_{\rm o}^*({\rm u})}{f_{\rm o}^2(\xi_{\rm p}^0)}$$

with $1 - F(u) = [1 - F_0(u)][1 - H(u)]$ and $F_0^*(u) = P(X_i \le u, \Delta_i = 1)$, the subdistribution function of the uncensored observations.

Note that an example of a bandwidth sequence that satisfies the conditions of Theorem 3.1 is $h_n = cn^{-\delta}$ with $\delta > 1/4$.

The next theorem gives some conditions for which Q_n^\star and Q_n are asymptotically uniformly equivalent in mean square.

Theorem 3.2. Suppose F_o and H are continuous and (h.1), (K.1), and (K.3) hold. Assume $E(X_1^{2q}) < \infty$ for some q > 1, where $X_1 = \min\{X_1^0, U_1\}$. Let η be such that $[1-F_o(\eta)][1-H(\eta)] > 0$ and let $T^* = F_o(\eta)$. Then for all $T \in [0,T^*)$, $\lim_{n\to\infty} E[\sup_{0\le p\le T} |Q_n^*(p) - Q_n(p)|^2] = 0$, provided $n^{1/2}h_n^4 \to \infty$ as $n \to \infty$.

4. Proofs of Theorems

The following two lemmas will be needed in the proof of Theorem 3.1.

In this section, $\{K(s,t): 0 \le s \le 1, t \ge 0\}$ will denote the generalized Kiefer process as stated by Csörgó (1983, Ch. 8).

<u>Lemma 1</u>. For 0 , where <math>T < 1, and $\delta < \min\{T - p, p\}$, as $n \rightarrow \infty$

 $\sup_{|h|<\delta} |n^{-1/2}[K(p+h,n) - K(p,n)]| \to 0 \text{ in probability.}$

The proof of Lemma 1 follows the same argument as the proof of Theorem 8.2.1 of Csorgo (1983).

Lemma 2. Suppose the derivative f_0' is continuous at ξ_p^0 and $f_0(\xi_p^0) > 0$. Under assumptions (h.1), (K.1), (K.2), (F.1) and (F.2), for 0 ,

$$\left| \int_0^1 \left[q_n(t) - q_n(p) \right] h_n^{-1} K\left(\frac{t-p}{h_n}\right) dt \right| \to 0$$
 in probability

as $n \to \infty$, where $q_n(t) = n^{1/2} [\hat{Q}_n(t) - Q^0(t)]$ denotes the product-limit quantile process.

<u>Proof.</u> For any given $\delta > 0$, there exists N such that when $n \ge N$,

$$|f_{0}^{1}[q_{n}(t) - q_{n}(p)] h_{n}^{-1} K(\frac{t-p}{h_{n}})dt|$$

$$= |f_{A(\delta)}[q_{n}(t) - q_{n}(p)] h_{n}^{-1} K(\frac{t-p}{h_{n}})dt|$$

$$\leq \sup_{t \in A(\delta)} |q_{n}(t) - q_{n}(p)|, \qquad (4.1)$$

where $A(\delta) = [p-\delta, p+\delta]$. By the conditions on f_0 , for δ sufficiently small, $f_0(Q^0(t)) > 0$ for all $t \in A(\delta)$. Hence, the right-hand-side of (4.1) is less than or equal to

$$\sup_{t \in A(\delta)} |\beta_{n}(t) - \beta_{n}(p)| \cdot |\frac{1}{f_{o}(Q^{o}(t))}| + \sup_{t \in A(\delta)} |\beta_{n}(p)[\frac{1}{f_{o}(Q^{o}(t))} - \frac{1}{f_{o}(Q^{o}(p))}]|,$$

where $\tilde{\rho}_n(t) = f_o(Q^o(t))q_n(t)$.

Let

$$a = \sup_{t \in A(\delta)} \left| \frac{1}{f_0(Q^0(t))} \right|$$
 and $b = \sup_{t \in A(\delta)} \left| \frac{1}{f_0(Q^0(t))} - \frac{1}{f_0(Q^0(p))} \right|$.

From Corollary 1 of Cheng (1981), since f_o is continuous at ξ_p^o , $\tilde{\rho}_n(p) + Z$ in distribution as $n + \infty$, where Z is a normally distributed random variable with mean zero and variance $\sigma^2 = (1-p)^2 \int_0^{\xi_p^o} [1-F(u)]^{-2} dF_o^*(u)$, with $1-F(u) = [1-F_o(u)][1-H(u)]$ and $F_o^*(u) = P(X_i \le u, \Delta_i = 1)$. Therefore,

$$\sup_{t \in A(\delta)} \left| \tilde{\rho}_{n}(p) \left[\frac{1}{f_{o}(Q^{o}(t))} - \frac{1}{f_{o}(Q^{o}(p))} \right] \right| \leq b \left| \tilde{\rho}_{n}(p) \right| \tag{4.2}$$

and for given $\varepsilon > 0$

$$\lim_{n\to\infty} P(|\tilde{\rho}_n(p)| > \varepsilon/b) \le P(|Z| > \varepsilon/b) \le b\sigma^2/\varepsilon^2. \tag{4.3}$$

Now.

$$\sup_{t \in A(\delta)} |\tilde{\rho}_{n}(t) - \tilde{\rho}_{n}(p)| \cdot \left| \frac{1}{f_{o}(Q^{o}(t))} \right| \leq a \sup_{t \in A(\delta)} |\tilde{\rho}_{n}(t) - \tilde{\rho}_{n}(p)|$$

$$\leq a \{ \sup_{t \in A(\delta)} |\tilde{\rho}_{n}(t) - n^{-1/2} |K(t,n)| + \sup_{t \in A(\delta)} |\tilde{\rho}_{n}(p) - n^{-1/2} |K(p,n)| + \sup_{t \in A(\delta)} |n^{-1/2} |K(p,n) - K(t,n)| \right\}. \tag{4.4}$$

For small enough δ , $p + \delta < T < 1$ and $p - \delta \ge 0$, so that by Corollary 8.3.3 of Csörgő (1983) as $n \to \infty$

$$\sup_{t \in A(\delta)} |\beta_n(t) - n^{-1/2} K(t,n)| \to 0 \text{ in probability}$$

and

$$\sup_{t\in A(\delta)} \left| \tilde{\rho}_n(p) - n^{-1/2} K(p,n) \right| \to 0 \quad \text{in probability.}$$

By Lemma 8.2.1 of Csörgo (1983) (or Berkes and Philipp, 1977) with $0 = t_0 < t_1 = n$, letting B(s) $\equiv n^{-1/2} K(s,n)$, $0 \le s \le 1$,

$$\sup_{t \in A(\delta)} |B(p) - B(t)| = \sup_{h \mid <\delta} |B(p+h) - B(p)|,$$

which by Lemma 1 converges to zero in probability for sufficiently small δ . Therefore, (4.4) converges to zero in probability as $n \to \infty$.

Finally, since b depends on δ , letting b become arbitrarily small gives from (4.2) and (4.4) that

$$\sup_{t \in A(\delta)} |\beta_n(p)[\frac{1}{f_o(Q^o(t))} - \frac{1}{f_o(Q^o(p))}]| \to 0 \text{ in probability.}$$

Thus, the result follows. ///

<u>Proof of Theorem 3.1.</u> Analogous to the beginning of the proof of Theorem 1 of Yang (1985), write

$$\sqrt{n} \left[Q_{n}(p) - Q^{0}(p) \right] = \int_{0}^{1} \left[q_{n}(t) - q_{n}(p) \right] h_{n}^{-1} K(\frac{t-p}{h_{n}}) dt
+ n^{1/2} \left[\int_{0}^{1} Q^{0}(t) h_{n}^{-1} K(\frac{t-p}{h_{n}}) dt - Q^{0}(p) \right] + q_{n}(p), \quad (4.5)$$

where $q_n(t) = n^{1/2}[\hat{Q}_n(t) - Q^0(t)]$ as in Lemma 2.

From Lemma 2, the first term on the right hand side of (4.5) is o (1), which means that it converges to zero in probability as $n \to \infty$. Similar to Yang's (1985) equation (10),

$$n^{1/2} \left[\int_{0}^{1} Q^{O}(t) h_{n}^{-1} K(\frac{t-p}{h_{n}}) dt - Q^{O}(p) \right]$$

$$= o(n^{1/2} h_{n}^{2}) + n^{1/2} h_{n}^{2} Q^{O''}(p) \int_{-\infty}^{\infty} \frac{t^{2}}{2} K(t) dt. \tag{4.6}$$

With the assumption that $n^{1/4}h_n \to 0$ as $n \to \infty$, (4.6) is also $p^{0}(1)$. Therefore, by Corollary 1 of Cheng (1981), the conclusion of the theorem follows. ///

Proof of Theorem 3.2. For $0 \le p \le T$, write

$$Q_n^*(p) - Q_n(p) = h_n^{-1} \sum_{i=1}^n Z_i[s_i K(\frac{S_i^{-p}}{h_n}) - \int_{S_{i-1}}^S K(\frac{t-p}{h_n})dt].$$

When $s_i > 0$, that is, Z_i is uncensored, let S_i^* be an interior point of the interval (S_{i-1}, S_i) with probability one so that

$$s_i K((S_i^*-p)/h_n) = \int_{S_{i-1}}^{S_i} K((t-p)/h_n) dt$$
 almost surely.

Then by condition (K.3), letting $I_{\hat{A}}$ denote the indicator function of the set A,

$$\begin{split} |Q_{n}^{*}(p) - Q_{n}(p)|I_{[0,T]}(p) &\leq h_{n}^{-1} \sum_{i=1}^{n} s_{i}Z_{i}|K((S_{i}-p)/h_{n}) \\ &- K((S_{i}^{*}-p)/h_{n})|I_{[0,T]}(p) I_{[S_{i}^{*}-ch_{n},1]}(p) \\ &\leq \Gamma h_{n}^{-2} \sum_{i=1}^{n} Z_{i}s_{i}|S_{i}-S_{i}^{*}|I_{[0,T]}(p) I_{[S_{i}^{*}-ch_{n},1]}(p) \\ &\leq \Gamma h_{n}^{-2} \sum_{i=1}^{n} Z_{i}s_{i}^{2} I_{[0,T]}(p) I_{[S_{i}^{*}-ch_{n},1]}(p) \end{split}$$

So

$$|Q_{n}^{*}(p) - Q_{n}(p)|^{2} I_{[0,T]}(p) \leq \Gamma^{2}h_{n}^{-4}(\Sigma Z_{i}s_{i}^{2} I_{[0,T]}(p) I_{[S_{i}^{*}-ch_{n},1]}(p))^{2}.$$
 (4.7)

Now,

$$(\sum_{i} z_{i}^{2} I_{[0,T]}^{(p)} I_{[S_{i}^{*}-ch_{n},1]}^{(p)})^{2}$$

$$\leq \sum_{i} z_{i}^{2} I_{[0,T]}^{3} I_{[0,T]}^{(p)} I_{[S_{i}^{*}-ch_{n},1]}^{(p)}$$

$$\leq \sum_{i} z_{i-1}^{2} (|\hat{F}_{n}(z_{i}) - F_{o}(z_{i})| + |F_{o}(z_{i}^{*}) - \hat{F}_{n}(z_{i}^{*})|)^{3} I_{[0,T]}^{(p)} I_{[S_{i}-2^{-ch_{n},1}]}^{(p)}, (4.8)$$

where g(x) denotes the limit from the left at x of the function g.

There is an N such that for all $n \ge N$, $T + ch_N < T^*$ and $T_N < F_O(\eta - \epsilon)$ where $T_N = T + ch_N$. For all such n's

$$\sum_{i=1}^{2} (|\hat{\mathbf{f}}_{n}(z_{i}) - \mathbf{f}_{o}(z_{i})| + |\mathbf{f}_{o}(z_{i}^{-}) - \hat{\mathbf{f}}_{n}(z_{i}^{-})|)^{3} \mathbf{I}_{[0,T]}(p) \mathbf{I}_{[S_{i-2}-ch_{n},1]}(p)$$

$$\leq 8 \sum_{i=1}^{2} \sup_{0 \leq x \leq \hat{\mathbf{Q}}_{n}(T_{N})} |\hat{\mathbf{f}}_{n}(x) - \mathbf{f}_{o}(x)|^{3},$$

$$(4.9)$$

which is independent of p. Notice also that

$$\sup_{0\leq x\leq \hat{Q}_n(T_N)+\epsilon} \left|\hat{F}_n(x)-F_o(x)\right|^3 \leq \sup_{0\leq x\leq \eta} \left|\hat{F}_n(x)-F_o(x)\right|^3 + \mathbb{I}_{\left[\hat{Q}_n(T_N)>\eta-\epsilon\right]}.$$

From the exponential bound in Theorem 2 of Földes and Rejtö (1981), for any $r \ge 1, \quad \{n^{1/2} \sup_{0 \le x \le \eta} |\hat{F}_n(x) - F_o(x)|\}^r, \quad n \ge 1\} \quad \text{is uniformly integrable.} \quad \text{Also,}$

$$\begin{split} & E(n^{3/2} \ I_{\{\hat{Q}_{n}(T_{N}) > \eta - \epsilon\}})^{r} \\ & = n^{3r/2} P[\hat{Q}_{n}(T_{N}) > \eta - \epsilon] \\ & = n^{3r/2} P[\hat{Q}_{n}(T_{N}) - F_{o}^{-1}(T_{N}) > \eta - F_{o}^{-1}(T_{N}) - \epsilon]. \end{split}$$

Letting $\gamma = \eta - F_0^{-1}(T_N) - \epsilon$, where ϵ is chosen so that $\gamma > 0$,

$$\begin{split} \mathbb{P} [\hat{Q}_{\mathbf{n}}(T_{\mathbf{N}}) &- F_{\mathbf{o}}^{-1}(T_{\mathbf{N}}) > \gamma] \\ & \leq \mathbb{P} [T_{\mathbf{N}} > \hat{F}_{\mathbf{n}}(\eta - \varepsilon)] \\ & = \mathbb{P} [T_{\mathbf{N}} - F_{\mathbf{o}}(\eta - \varepsilon) > \hat{F}_{\mathbf{n}}(\eta - \varepsilon) - F_{\mathbf{o}}(\eta - \varepsilon)] \\ & \leq \mathbb{P} [|\hat{F}_{\mathbf{n}}(\eta - \varepsilon) - F_{\mathbf{o}}(\eta - \varepsilon)| > F_{\mathbf{o}}(\eta - \varepsilon) - T_{\mathbf{N}}], \end{split}$$

By the same exponential bound as in Theorem 2 of Földes and Rejtö (1981), as $n \rightarrow \infty$

$$\mathbb{E}(n^{3/2} \mathbb{I}_{\left[\hat{Q}_{n}(T_{N}) > \eta - \varepsilon\right]})^{r} \to 0.$$

Therefore, $\{(n^{1/2} \sup_{0 \le x \le \hat{Q}_n(T_N) + \epsilon} |\hat{F}_n(x) - F_o(x)|\}^r$, $n \ge 1$ is uniformly integrable.

By hypothesis, $E(X_1^{2q}) < \infty$ for some q > 1, so $\{(n^{-1} \sum_{i=1}^{n} z_i^2)^q, n \ge 1\}$ is uniformly integrable. Thus,

for
$$\frac{1}{s} + \frac{1}{q} = 1$$
,

$$E(n^{3/2} \sup_{0 \le x \le \hat{Q}_n(T_n) + \varepsilon} |\hat{F}_n(x) - F_o(x)|^3 n^{-1} \sum_{i=1}^n z_i^2)$$

$$\leq (\mathbb{E}(n^{3/2} \sup_{0 \leq x \leq \hat{Q}_{n}(T_{N}) + \epsilon} |\hat{F}_{n}(x) - F_{0}(x)|^{3})^{s})^{1/s} \mathbb{E}((n^{-1} \sum_{i=1}^{n} z_{i}^{2})^{q})^{1/q}$$

Therefore, from (4.7) - (4.10), by the hypothesis $n^{1/2}h_n^4 \rightarrow \infty$,

$$E[\sup_{0 \le p \le T} |Q_n^*(p) - Q_n(p)|^2] = o(1),$$

completing the proof. ///

References

で人に人のないが、このではないとうと

- Berkes, I. and Philipp, W. (1977). An almost sure invariance principle for the empirical distribution function of mixing random variables. Z. Wahrsch. Verw. Gebiete 41, 115-137.
- Cheng, K. F. (1981). On almost sure representations for quantiles of the product limit estimator with applications. Research Report No. 87, SUNY Buffalo, Dept. of Statistics.
- Csörgó, M. (1983). Quantile Processes with Statistical Applications. CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, PA.
- Efron, B. (1967). The two-sample problem with censored data. Proc. Fifth Berkeley Sympos. 4, 831-853.
- Falk, M. (1984). Relative deficiency of kernel type estimators of quantiles.

 Ann. Statist. 12, 261-268.
- Falk, M. (1985). Asymptotic normality of the kernel quantile estimator. Ann. Statist. 13, 428-433.
- Földes, A. and Rejtö, L. (1981). A LIL type result for the product limit estimator. Z. Wahr. Verw. Gebiete 56, 75-86.
- Kaplan, E. L. and Meier, P. (1958). Nonparametric estimation from incomplete observations. J. Amer. Statist. Assoc. 53, 457-481.
- Padgett, W. J. (1985). A kernel type estimator of a quantile function from right-censored data. Statistics Technical Report No. 101R, University of South Carolina, Dept. of Math. and Statist.
- Parzen, E. (1979). Nonparametric statistical data modeling. <u>J. Amer. Statist.</u>
 <u>Assoc. 74</u>, 105-121.
- Sander, J. (1975). The weak convergence of quantiles of the product limit estimator. Technical Report 5, Stanford University, Dept. of Statistics.
- Yang, S. S. (1985). A smooth nonparametric estimator of a quantile function. Technical Report, Kansas State University, Dept. of Statistics.

BECURITY CLASSIFICATION OF THIS PAGE

SECURITY CLASSIFICATION OF THIS PAGE								
REPORT DOCUMENTATION PAGE								
14 REPORT SECURITY CLASSIFICATION UNCLASSIFIED	16. NESTRUCEIVE MILLO 302							
24 SECURITY CLASSIFICATION AUTHORITY	3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution							
26. DECLASSIFICATION/DOWNGRADING SCHED	unlimited.							
4. PERFORMING ORGANIZATION REPORT NUM	5. MONITORING ORGANIZATION REPORT NUMBER(8)							
Stat. Tech. Rep. No. 104 (62G0	AFOSR-TR- 5- 865							
6a NAME OF PERFORMING ORGANIZATION Department of Mathematics	Bb. OFFICE SYMBOL (If applicable)	7& NAME OF MONITORING ORGANIZATION			:			
and Statistics		Air Force Office of Scientific Research			earch			
6c. ADDRESS (City, State and ZIP Code)	·	7b. ADDRESS (City, State and ZIP Code)						
University of South Carolina	••	Directorate of Mathematical & Information			rmation			
Columbia SC 29208		Sciences, Bolling AFB DC 20332						
84. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER						
AFOSR , ARO	NM	AFOSR-84-0156						
Sc. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NOS.						
Bolling AFB DC 20332	÷	PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT			
		61102F	2304	A5	NO.			
13. TITLE (Include Security Classification)								
ON THE ASYMPTOTIC PROPERTIES OF A KERNEL TYPE QUANTILE ESTIMATOR FROM CENSORED SAME								
12. PERSONAL AUTHOR(S) Y. L. Lio, W. J. Padgett, and K. F. Yu								
Technical 136 TIME CO	OVERED TO	14. DATE OF REPORT (Yr., No., Dey) 15. PAGE COUNT 1985 May 13						
16. SUPPLEMENTARY NOTATION								
17. COSATI CODES 18. SUBJECT TERMS (Continue on reserve if necessary and identify by block number)								
FIELD GROUP SUB. GR.		censorship; Kernel estimation; Product-limit						
	quantile func							
	vergence.							
19. ASSTRACT (Continue on reverse if necessary and identify by block number) Some asymptotic results for a kernel type estimator of the quantile function from right-								
			9 4 .		_			
censored data are obtained. The estimator is defined by $Q_n(p) = h_n^{-1} \int_0^1 \hat{Q}_n(t) R((t-p)/h_n) dt$,								
which is smoother than the usual product-limit quantile function $\hat{Q}_n(p)=\inf\{t: \hat{F}_n(t) \geq p\}$,								
where \hat{F}_n denotes the product-limit estimator of the lifetime distribution F_0 . Under the								
random censorship model and general conditions on h_n , K , and F_0 , the asymptotic normality								
of $Q_{\mathbf{n}}(\mathbf{p})$ is proven. In addition, as approximation to $Q_{\mathbf{n}}$ is shown to be asymptotically								
uniformly equivalent to Q_n in mean square.								
•					f			
i i								
20. DISTRIBUTION/AVAILABILITY OF ASSTRAC	21. ABSTRACT SECURITY CLASSIFICATION							
UNCLASSIFIED/UNLIMITED 🖾 SAME AS RPT.	UNCLASSIFIED							
224 NAME OF RESPONSIBLE INDIVIDUAL	22b. TELEPHONE NUMBER 22c. OFFICE SYMBOL Anclude Area Code;							
CPT Brian W. Woodruff	(202) 767-502		NM					
DO FORM 1472 CO 422								

END

FILMED

11-85

DTIC